

STUDIES OF MEXICAN AND CENTRAL AMERICAN
SPECIES OF SENEPIO

J. M. Greenman¹

SENECIO ANDRIEUXII DC., Prodr. 6: 430. 1837; Hemsl.
in Biol. Cent.-Am. Bot. 2: 236. 1881.

MEXICO: Sierra Madre, Michoacán and Guerrero, alt.
2000 m., Feb. 1899, *Langlassé* 895 (Herb. De Candolle,
Geneva).

SENECIO ASCHENBORNIANUS Schauer in Linnaea 20: 698.
1847.

MEXICO: Chiefly in the region of San Luis Potosí, 22°
N. latitude, alt. 1800-2400 m., 1878, *Parry & Palmer* 539
(Phil. Acad. Sci.); hills near Jalapa, Vera Cruz, alt. 4000
m. April 1, 1899, *Pringle* 8069 (Phil. Acad. Sci. and Mo.
Bot. Gard.); Chontla, Vera Cruz, alt. 508 m., March 1910,
Cárdenas 497 and 521 (Mo. Bot. Gard.); Sierra San Lucas,
Jaumave, Tamaulipas, June 1933, *von Rozynski* 654 and 693
(Mo. Bot. Gard.); on moist open pine-oak slopes, Dul-
ces Nombres, Nuevo León and just east of the border into
Tamaulipas, alt. 1690 m., July 18, 1948, *Meyer & Rogers*
2800 (Mo. Bot. Gard.).

Senecio Bartlettii *Greenm. sp. nov.*

Herbaceus perennis, juventate ubique plus minusve floc-
coso-tomentulosus; caule simplici vel basi ramoso, 4.5-6 dm.
alto, aliquanto glabratu; foliis pinnato-lobatis, plus minusve
petiolatis, amplexicaulibus, supra arachnoideo-tomentulosis,
subtus denique persistenter albo-tomentosis; lobis lateralibus
oblongis, acutis vel obtusis, lobo terminali rhombeo-ovato vel
reniformi; inflorescentiis corymbo-cymosis; capitulis ca. 1 cm.
altis; involucris campanulatis, minute bracteolatis; involucri
reniformi; inflorescentiis corymbo-cymosis; capitulis ca. 1 cm.

¹ Curator Emeritus of the Herbarium of the Missouri Botanical Garden; Professor Emeritus in the Henry Shaw School of Botany of Washington University.

longis; floribus femineis ligulatis, ligulis flavis; floribus disci ca. 60; achaeniis maturitate striatis, hirtellis.

MEXICO: Mesa de Tierra, vicinity of San José, Tamaulipas, alt. 1060 m., July 19, 1930, Bartlett 10427 (type in Univ. Mich. Herb.); Cerro Barril, vicinity of San José, Taumalipas, alt. 1400 m., July 19, 1930, Bartlett 10422a (Univ. Mich. Herb.).

Senecio cobanensis Coulter var. *sublaciniatus*
Greenm., var. nov.

Frutex, 2-5 m. altus; foliis petiolatis, alternis, elliptico-lanceolatis, 6-17 cm. longis, 1-4 cm. latis, utrinque acuminatis, acutis, subintegris, sinuato-denticulatis vel laciniato-dentatis, utrinque glabris; petiolis 1-6 cm. longis; inflorescentiis juv-entate albo-tomentulosis, glabratis; involucri squamis 5; floribus 4-5, flavis, 1 vel 2 ligulatis; achaeniis glabris.

GUATEMALA: in damp forest, hills of finca Carmona southeast of Antigua, Sacatépequez, shrub or tree 2-5 m., alt. 1590-1800 m., Jan. 27, 1939, Standley 63680 (Mo. Bot. Gard. Herb., TYPE; Chicago Nat. Hist. Mus. Herb.); Volcán Sta. Clara, south facing slopes to the summit, alt. 2100-3000 m., June 5, 1942, Steyermark 46994 (Mo. Bot. Gard. and Chicago Nat. Hist. Mus.).

SENECIO HARTWEGII Benth., Pl. Hartw. 18. 1839.

MEXICO: trail from El Batel to Pico del Aguila, Sierra Madre, Sinaloa, alt. 1200 m., Nov. 11, 1925, Mexia 441 (Copenhagen Herb.).

SENECIO HETERODONTUS Greenm. in Field Columb. Mus. Bot. 2: 277. 1907.

MEXICO: Alvarez, San Luis Potosí, July 13-23, 1904, Palmer 237 (Copenhagen Herb.).

The specimen, namely Palmer 237 in the United States National Herbarium, is somewhat intermediate between *S. heterodontus* Greenm. and *S. multidentatus* Schz. Bip. ex Hemsley.

SENECIO HOFFMANII Klatt in Leopoldina **25:** 106. 1899
(reprint p. 3).

MEXICO: Tapachula, Chiapas, alt. 90 m., Jan. 27, 1911,
Quarles van Uffort 79 (Bot. Mus. Utrecht).

SENECIO JURGENSENII Hemsl., Biol. Cent.-Am. Bot. **2:**
242. 1881.

MEXICO: rocky shaded slope, open forest, Cerro Tiburón,
Sierra Tacuichamona, Sinaloa, alt. 1400 m., Feb. 18, 1940,
Gentry 5659 (Univ. Mich. and Mo. Bot. Gard.).

Senecio lithophilus Greenm., sp. nov.

Herbaceus perennis; caule erecto vel adscendente
simplice vel ramoso 40-60 cm. alto, albo-tomentoso, usque
ad inflorescentiam folioso; foliis inferioribus oblanceolatis 5-
20 cm. longis, 1-4 cm. latis aliquanto et irregulariter sinuato-
denticulatis, caulinis sessilibus semiamplexicaulibus, oblongo-
lanceolatis, 5-20 cm. longis, 1-4 cm. latis, supra juventate flo-
ccoso-tomentulosis mox glabratis, subtus persistenter albo-tomen-
tosis; inflorescentiis terminalibus corymboso-cymosis, capitulis
in anthesi 10-12 mm. altis, heterogamis; involucri bracteis
21-25, linear-lanceolatis, acutis, sparse floccoso-tomentulosis,
plus minusve glabratis; floribus liguliferis circiter 18, ligulis
anguste oblongo-linearibus 10-15 mm. longis, flavis, tubo
gracili circiter 5 mm. longo, floribus disci numerosis, circiter
80; achaeniis sparse hirtellis.

MEXICO: on open faces of limestone cliffs on road from
Adelaida, Tamaulipas to Dulces Nombres, Nuevo León but in
the state of Tamaulipas, alt. 1690 m., July 12, 1948, Meyer
& Rogers 2751 (TYPE in Mo. Bot. Gard. Herb.).

Senecio lithophilus is related to *S. huachucanus* Gray, but
differs in being copiously tomentose, and in having larger
heads and more numerous flowers.

SENECIO LORATIFOLIUS Greenm. in Ann. Mo. Bot. Gard.
5: 93. 1948.

MEXICO: in mountain meadows 6 kilometers west of Miquihuana, Tamaulipas, alt. 2940 m., August 6, 1941, *Stanford, Rutherford, and Northcraft* 734 (Mo. Bot. Gard.); densely tufted perennial to 40 cm. tall, in open pine forest, Cerro del Viejo, 24 kms. west of Dulces Nombres, municipality of Zaragoza, Nuevo León, alt. 3330 m., August 18, 1948, *Meyer & Rogers* 2979 (Mo. Bot. Gard.).

Senecio morazensis Greenm., sp. nov.

Frutex scandens ubique glaber, caule tereti in sicco cortice brunneo tecto; foliis alternis, plus minusve secundis, petiolatis, elliptico-lanceolatis, 4-8 cm. longis, 1.5-3 cm. latis, acuminatis, acutis, integris, basi cuneatis; petiolis usque ad 10-12 mm. longis; inflorescentiis paniculatis; capitulis heterogamis ca. 1 cm. longis; involucris sparse calyculatis, campanulatis; involucri squamis 8, lineari-lanceolatis vel oblongis, 6-7 mm. longis, usque ad 4 mm. latis; floribus femineis 8 ligulatis, ligulis anguste lineari-oblongis, 5-6 mm. longis; floribus disci circiter 30; achaeniis glabris.

HONDURAS: flowers yellow, vine in tree in cloud forest area in mountains above San Juancito, Morazán, alt. 2000 m., March 25, 1948, *Williams & Molina* 13976 (TYPE in Mo. Bot. Gard. and ISOTYPE in Herb. Escuela Agrícola Panamericana); liana de 1-3 m., floresta de nubes de la montaña La Tigra, suroeste de San Juancito, Morazán, 5 febrero 1950, *Williams & Molina* 17104 (Herb. Escuela Agrícola Panamericana).

Senecio morazensis belongs to the section *Streptothamni* and it is allied to *S. candelariae* Benth. and *S. parasiticus* Hemsl. In general habit, stem and leaves, there is almost complete accord with *S. parasiticus*, but *S. morazensis* differs in having heterogamous heads with larger diameter, longer and wider involucral bracts with wide margin.

SENECIO MULGEDIIFOLIUS Schauer in Linnaea **19**: 733. 1847; Hemsl. Biol. Cent.-Am. Bot. **2**: 243. 1881; Greenm. Monog. Senecio, pt. I, p. 25 and in Engler's Bot. Jahrb. **32**: 21. 1902.

MEXICO: S. Miguel el Soldado, Vera Cruz, June-Oct. 1840, *Galeotti* 2291 (Bot. Gard., Brussels); bosque de encinos, Angostura, D. F., alt. 2600 m., Sept. 1941, *Lyonnet* 3427 (U. S. Nat. Herb.).

SENEPIO MULTIDENTATUS Schz. Bip. ex Hemsl., Biol. Cent.-Am. Bot. 2: 243. 1881.

MEXICO: by streams, Sierra de las Cruces, Mexico, alt. 3000 m., Aug. 28, 1904, *Pringle* 13077 (Copenhagen Herb.); Vaqueria del Jacal, alt. 2500-3200 m., June-Oct. 1840, *Galeotti* 2164 (Bot. Gard., Brussels); by streams, Sierra de las Cruces, México, Aug. 23, 1892, *Pringle* 4205 (Phil. Acad. Nat. Sci., Mo. Bot. Gard. and Vienna Herb.); by streams, Sierra de las Cruces, Mexico, alt. 3000 m., Sept. 3, 1901, *Pringle* 9314 (Mo. Bot. Gard.); orilla del arroyo Bosque, Salazar, (Mexico), alt. 2900 m., Sept. 1941, *Lyonnet* 3375 (U. S. Nat. Herb.); La Cima, alt. 3000 m., Aug. 1904, *Kuntze* s. n. (N. Y. Bot. Gard.).

SENEPIO PHANERANDRUS Cufodontis in Arch. Bot., Forli 9: 103. 1933.

COSTA RICA: Volcán Poás, alt. 2400 m., April 26, 1930, *Cufodontis* 544 (Mo. Bot. Gard.); Volcán Poás, alt. 2678 m., January 30, 1922, J. M. & M. T. Greenman 5392 (Mo. Bot. Gard.); Volcán Poás, alt. 2500-2575 m., Dec. 1-Jan. 1, 1938, *Allen* 616 (Mo. Bot. Gard.); Volcán Poás, alt. 2600 m., March 31, 1907 *Pittier* 2050 (U. S. Nat. Herb.); Volcán Poás, alt. 2500-2640 m., Feb. 17, 1924, *Standley* 34840 (U. S. Nat. Herb.); environs de la Lagune du Barba, alt. 2755, Feb. 6, 1890, "A. C.", 1894 (U. S. Nat. Herb.).

SENEPIO QUEBRADENSIS Greenm. in Ann. Mo Bot. Gard. 3: 117. 1916.

MEXICO: paraje de los Sabinos, San Ignacio, Sinaloa, alt. 510 m., March 9, 1918, *Narvaez & Salazar* 267 (U. S. Nat. Herb.); steep clay slopes with brush in oak forest, Puerto a Tamiapa, Sinaloa, alt. 1400 m., March 6-8, 1940, *Gentry* 5825 (Mo. Bot. Gard.).