

Agrícola Panamericana, dando flores todos los años. He hecho averiguaciones en varias partes, resultando que esta forma es poco conocida en Honduras. No tengo datos de su existencia en otros países centroamericanos.

Búsqueda de la literatura hortícola disponible ha resultado en una sola referencia a Amaryllis de flores dobles: en el recién (1951) publicado "Dictionary of Gardening," de la Royal Horticultural Society de Londres, hay mención de una *Hippeastrum equestre*, "form flore pleno," sus flores anaranjado rico. Las flores de las plantas hondureñas son escarlatas o rojo anaranjado.


MASDEVALLIA BORUCANA, A NEW ORCHID FROM COSTA RICA

PAUL H. ALLEN

Masdevallia borucana P. H. Allen, sp. nov.

Folia coriacea, spatulata, apice obtusa vel retusa; scapi filiformes, uniflori; pars basalis sepolorum lateralium 2.25-2.50 cm. longa, usque ad 1.5 cm. lata, late subreniformis; petala ca. 6 mm. longa, ambitu oblongo-pandurata; labium carnosum, lamina ca. 5 mm. longa et 2 mm. lata.

Caespitose epiphytic herbs, to about 15 cm. in height. Secondary stems very short, 5-10 mm. long. Leaves coriaceous, spatulate in outline, 8-14 cm. long and 1.5-2.5 cm. wide, with obtuse or retuse apices, the apical notch usually with a minute but definite central apicule. Scapes filiform, 1-flowered, about equalling the leaves in length. Sepaline tube somewhat inflated, composed of very unequal parts and strongly gibbose in profile; dorsal sepal joined to the laterals for only about 1 cm., the upper free portion being produced into a filiform, erect cauda 5-6 cm. long; basal portion of the lateral sepals 2.25-2.5 cm. long and up to 1.5 cm. wide, broadly subreniform in outline and connate for about 2/3 of their length, the convex terminal portions abruptly contracted into filiform appendages 3-3.5 cm. long. Petals about 6 mm. long and 2 mm. wide, from a narrowed base oblong-pandurate in outline, the outer surface minutely papillose and obscurely sulcate near the broadly truncate apex, the inner surface provided with a



Masdevallia borucana. Fig. 1, petals, column and lip seen in profile. Fig. 2, the lip. Fig. 3, front view of the column and petals, showing undulate keels.

longitudinal undulate keel, terminating in a short point on the outer lobe of the apex of each petal. The fleshy lip is articulated with the foot of the column, the minutely papillose, ligulate blade about 5 mm. long and 2 mm. wide, the apical 1/3 somewhat narrowed and reflexed in natural position. Column about equalling the petals in length, the margins of the hooded apex irregularly and minutely serrate.